



Cambridge International AS & A Level

PHYSICS

9702/23

Paper 2 AS Level Structured Questions

October/November 2021

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2021 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **12** printed pages.

Abbreviations

/	Alternative and acceptable answers for the same marking point.
()	Bracketed content indicates words which do not need to be explicitly seen to gain credit but which indicate the context for an answer. The context does not need to be seen but if a context is given that is incorrect then the mark should not be awarded.
—	Underlined content must be present in answer to award the mark. This means either the exact word or another word that has the same technical meaning.

Mark categories

B marks	These are <u>independent</u> marks, which do not depend on other marks. For a B mark to be awarded, the point to which it refers must be seen specifically in the candidate's answer.
M marks	These are <u>method</u> marks upon which A marks later depend. For an M mark to be awarded, the point to which it refers must be seen specifically in the candidate's answer. If a candidate is not awarded an M mark, then the later A mark cannot be awarded either.
C marks	These are <u>compensatory</u> marks which can be awarded even if the points to which they refer are not written down by the candidate, providing subsequent working gives evidence that they must have known them. For example, if an equation carries a C mark and the candidate does not write down the actual equation but does correct working which shows the candidate knew the equation, then the C mark is awarded. If a correct answer is given to a numerical question, all of the preceding C marks are awarded automatically. It is only necessary to consider each of the C marks in turn when the numerical answer is not correct.
A marks	These are <u>answer</u> marks. They may depend on an M mark or allow a C mark to be awarded by implication.

Annotations

✓	Indicates the point at which a mark has been awarded.
X	Indicates an incorrect answer or a point at which a decision is made not to award a mark.
XP	Indicates a physically incorrect equation ('incorrect physics'). No credit is given for substitution, or subsequent arithmetic, in a physically incorrect equation.

ECF	Indicates 'error carried forward'. Answers to later numerical questions can always be awarded up to full credit provided they are consistent with earlier incorrect answers. <u>Within</u> a section of a numerical question, ECF can be given after AE, TE and POT errors, but not after XP.
AE	Indicates an arithmetic error. Do not allow the mark where the error occurs. Then follow through the working/calculation giving full subsequent ECF if there are no further errors.
POT	Indicates a power of ten error. Do not allow the mark where the error occurs. Then follow through the working/calculation giving full subsequent ECF if there are no further errors.
TE	Indicates incorrect transcription of the correct data from the question, a graph, data sheet or a previous answer. For example, the value of 1.6×10^{-19} has been written down as 6.1×10^{-19} or 1.6×10^{19} . Do not allow the mark where the error occurs. Then follow through the working/calculation giving full subsequent ECF if there are no further errors.
SF	Indicates that the correct answer is seen in the working but the final answer is incorrect as it is expressed to too few significant figures.
BOD	Indicates that a mark is awarded where the candidate provides an answer that is not totally satisfactory, but the examiner feels that sufficient work has been done ('benefit of doubt').
CON	Indicates that a response is contradictory.
I	Indicates parts of a response that have been seen but disregarded as irrelevant.
M0	Indicates where an A category mark has not been awarded due to the M category mark upon which it depends not having previously been awarded.
^	Indicates where more is needed for a mark to be awarded (what is written is not wrong, but not enough). May also be used to annotate a response space that has been left completely blank.
SEEN	Indicates that a page has been seen.

Question	Answer	Marks
1(a)	$m = \rho V$ or ρAL	C1
	$W = mg$	C1
	$(A =) 24 / (9.81 \times 850 \times 0.18) = 0.016 \text{ (m}^2\text{)}$	A1
	or	
	$P = F / A$	(C1)
	$P = \rho gh$	(C1)
1(b)(i)	$(A =) 24 / (9.81 \times 850 \times 0.18) = 0.016 \text{ (m}^2\text{)}$	(A1)
	$(\text{upthrust} =) 24 + 8(.0) = 32 \text{ (N)}$	A1
1(b)(ii)	$(\Delta)p = 32 / 0.016 (= 2000)$	C1
	$(\Delta)p = \rho g(\Delta)h$	C1
	$\rho = 2000 / (9.81 \times 0.17)$	
	$= 1200 \text{ kg m}^{-3}$	A1
1(c)(i)	$E = \frac{1}{2}Fx$ or $E = \frac{1}{2}kx^2$ or $E = \text{area under graph}$	C1
	$(\Delta)E = (\frac{1}{2} \times 8.0 \times 0.40) - (\frac{1}{2} \times 4.0 \times 0.20)$	C1
	or	
	$(\frac{1}{2} \times 20 \times 0.40^2) - (\frac{1}{2} \times 20 \times 0.20^2)$	
	or	
1(c)(ii)	$\frac{1}{2} \times (4.0 + 8.0) \times 0.20$	
	$= 1.2 \text{ J}$	A1
1(c)(ii)	length = 30 cm	A1

Question	Answer	Marks
2(a)	force \times displacement in the direction of the force	B1
2(b)	units: $\text{kg m s}^{-2} \times \text{m} = \text{kg m}^2 \text{s}^{-2}$	A1
2(c)	$T_1: \text{K}$ and $T_2: \text{K}$	C1
	$A: \text{m}^2$ and $t: \text{s}$ and $L: \text{m}$	C1
	$c = (\text{kg m}^2 \text{s}^{-2} \text{m}) / (\text{m}^2 \text{K s})$ $= \text{kg m s}^{-3} \text{K}^{-1}$	A1

Question	Answer	Marks
3(a)	change in displacement / time (taken)	B1
3(b)	<p>by calculation: $v^2 = 42^2 + 23^2 - (2 \times 42 \times 23 \times \cos 54^\circ)$</p> <p>or $v^2 = (42 - 23 \cos 54^\circ)^2 + (23 \sin 54^\circ)^2$</p> <p>or $v^2 = (42 - 23 \sin 36^\circ)^2 + (23 \cos 36^\circ)^2$</p> <p>$v = 34 \text{ m s}^{-1}$</p> <p>or</p> <p>by scale diagram: triangle of vector velocities drawn</p> <p>$v = 34 \text{ m s}^{-1}$ (allow $\pm 1 \text{ m s}^{-1}$ if scale diagram used)</p>	C1
		A1
		(C1)
		(A1)
3(c)(i)	$(\Delta)E = mg(\Delta)h$ or $(\Delta)E = W(\Delta)h$ $h = 6100/46 (= 133 \text{ m})$ $\theta = \sin^{-1}(133/280)$ $= 28^\circ$	C1
		C1
		A1
3(c)(ii)	$\text{force} = 6100 / 280$ or $46 \sin 28^\circ$ $= 22 \text{ N}$	C1
		A1
3(d)	$v_{(s)} = 280 / 14 (= 20 \text{ m s}^{-1})$ $f_o = f_s v / (v - v_s)$ $f_s = 450 \times (340 - 20) / 340$ $= 420 \text{ Hz}$	C1
		C1
		A1

Question	Answer	Marks
4(a)	to the left/from the right/from B to A/opposite (direction) to (α -particle) velocity	B1
4(b)	$v^2 = u^2 + 2as$ $s = (4.1 \times 10^6)^2 / (2 \times 2.7 \times 10^{14})$ $= 0.031 \text{ m}$	C1 A1
4(c)	$E = F/Q$ or $E = ma/Q$ $= (4 \times 1.66 \times 10^{-27} \times 2.7 \times 10^{14}) / (2 \times 1.60 \times 10^{-19})$ $= 5.6 \times 10^6 \text{ V m}^{-1}$	C1 C1 A1
4(d)	straight line with negative gradient that intercepts both the momentum and t axes	B1
4(e)	force (on α -particle)	B1
4(f)(i)	$E = \frac{1}{2}mv^2$ $= \frac{1}{2} \times 9.11 \times 10^{-31} \times (4.1 \times 10^6)^2$ $= 7.7 \times 10^{-18} \text{ J}$	C1 C1 A1
4(f)(ii)	particles have opposite charges (so) forces (on charges) are opposite (directions) β^- has less/half the charge so less/half the force	B1 B1 B1
4(f)(iii)	(electron) <u>antineutrino</u>	B1

Question	Answer	Marks
5(a)	maximum displacement (of a point/particle on string/wave)	B1
5(b)	$v = \lambda / T$ or $v = f\lambda$ and $f = 1 / T$	C1
	$T = 690 \times 10^{-9} / 3.00 \times 10^8$	C1
	$= 2.3 \times 10^{-15} \text{ s}$	A1
5(c)(i)	$\lambda = ax / D$	C1
	$G = x / D$	A1
	(so) $a = \lambda / G$	
5(c)(ii)	straight line from origin always below printed line	M1
	line is half the height of printed line at maximum D	A1

Question	Answer	Marks
6(a)	$R = \rho L / A$ $(R =) 5.0 \times 10^{-7} \times 2.0 / 3.3 \times 10^{-7} = 3.0 \Omega$	C1 A1
6(b)(i)	$I = 1.2 / 3.0$ $= 0.40 \text{ A}$	A1
6(b)(ii)	$r = (1.50 - 1.20) / 0.40 \text{ or } 1.50 / 0.40 - 3.0$ $= 0.75 \Omega$	C1 A1
6(c)	$E / 1.20 = 1.4 / 2.0$ $E = 0.84 \text{ V}$ or $R_{XP} = (1.4 / 2.0) \times 3.0 (= 2.1 \Omega)$ $E = 2.1 \times 0.40$ $E = 0.84 \text{ V}$	C1 A1 (C1) (A1)
6(d)	(second wire has) larger resistance/resistance increases p.d. across XY is larger/increases (for second wire) or p.d. across the (second) wire is larger/increases (so) length XP (for second wire) is shorter	M1 M1 A1