



Cambridge International AS & A Level

PHYSICS

9702/22

Paper 2 AS Level Structured Questions

February/March 2022

MARK SCHEME

Maximum Mark: 60

<p>Published</p>

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the February/March 2022 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **12** printed pages.

PUBLISHED**Examples of how to apply the list rule**State **three** reasons.... [3]

A	1	Correct	✓	2
	2	Correct	✓	
	3	Wrong	✗	

B (4 responses)	1	Correct, Correct	✓, ✓	3
	2	Correct	✓	
	3	Wrong	ignore	

C (4 responses)	1	Correct	✓	2
	2	Correct, Wrong	✓, ✗	
	3	Correct	ignore	

D (4 responses)	1	Correct	✓	2
	2	Correct, CON (of 2.)	✗, (discount 2)	
	3	Correct	✓	

E (4 responses)	1	Correct	✓	3
	2	Correct	✓	
	3	Correct, Wrong	✓	

F (4 responses)	1	Correct	✓	2
	2	Correct	✓	
	3	Correct CON (of 3.)	✗ (discount 3)	

G (5 responses)	1	Correct	✓	3
	2	Correct	✓	
	3	Correct Correct CON (of 4.)	✓ ignore ignore	

H (4 responses)	1	Correct	✓	2
	2	Correct	✗	
	3	CON (of 2.) Correct	(discount 2) ✓	

I (4 responses)	1	Correct	✓	2
	2	Correct	✗	
	3	Correct CON (of 2.)	✓ (discount 2)	

Abbreviations

/	Alternative and acceptable answers for the same marking point.
()	Bracketed content indicates words which do not need to be explicitly seen to gain credit but which indicate the context for an answer. The context does not need to be seen but if a context is given that is incorrect then the mark should not be awarded.
—	Underlined content must be present in answer to award the mark. This means either the exact word or another word that has the same technical meaning.

Mark categories

B marks	These are <u>independent</u> marks, which do not depend on other marks. For a B mark to be awarded, the point to which it refers must be seen specifically in the candidate's answer.
M marks	These are <u>method</u> marks upon which A marks later depend. For an M mark to be awarded, the point to which it refers must be seen specifically in the candidate's answer. If a candidate is not awarded an M mark, then the later A mark cannot be awarded either.
C marks	These are <u>compensatory</u> marks which can be awarded even if the points to which they refer are not written down by the candidate, providing subsequent working gives evidence that they must have known them. For example, if an equation carries a C mark and the candidate does not write down the actual equation but does correct working which shows the candidate knew the equation, then the C mark is awarded. If a correct answer is given to a numerical question, all of the preceding C marks are awarded automatically. It is only necessary to consider each of the C marks in turn when the numerical answer is not correct.
A marks	These are <u>answer</u> marks. They may depend on an M mark or allow a C mark to be awarded by implication.

Annotations

✓	Indicates the point at which a mark has been awarded.
X	Indicates an incorrect answer or a point at which a decision is made not to award a mark.
XP	Indicates a physically incorrect equation ('incorrect physics'). No credit is given for substitution, or subsequent arithmetic, in a physically incorrect equation.
ECF	Indicates 'error carried forward'. Answers to later numerical questions can always be awarded up to full credit provided they are consistent with earlier incorrect answers. <u>Within</u> a section of a numerical question, ECF can be given after AE, TE and POT errors, but not after XP.

PUBLISHED

AE	Indicates an arithmetic error. Do not allow the mark where the error occurs. Then follow through the working/calculation giving full subsequent ECF if there are no further errors.
POT	Indicates a power of ten error. Do not allow the mark where the error occurs. Then follow through the working/calculation giving full subsequent ECF if there are no further errors.
TE	Indicates incorrect transcription of the correct data from the question, a graph, data sheet or a previous answer. For example, the value of 1.6×10^{-19} has been written down as 6.1×10^{-19} or 1.6×10^{19} . Do not allow the mark where the error occurs. Then follow through the working/calculation giving full subsequent ECF if there are no further errors.
SF	Indicates that the correct answer is seen in the working but the final answer is incorrect as it is expressed to too few significant figures.
BOD	Indicates that a mark is awarded where the candidate provides an answer that is not totally satisfactory, but the examiner feels that sufficient work has been done ('benefit of doubt').
CON	Indicates that a response is contradictory.
I	Indicates parts of a response that have been seen but disregarded as irrelevant.
M0	Indicates where an A category mark has not been awarded due to the M category mark upon which it depends not having previously been awarded.
^	Indicates where more is needed for a mark to be awarded (what is written is not wrong, but not enough). May also be used to annotate a response space that has been left completely blank.
SEEN	Indicates that a page has been seen.

PUBLISHED

Question	Answer	Marks
1(a)	F_v : kg m s^{-2}	C1
	k : $\text{kg m s}^{-2} / \text{m} \times \text{m s}^{-1}$ $= \text{kg m}^{-1} \text{s}^{-1}$	A1
1(b)	$F = \rho g V$ $V = 4/3 \times \pi \times (2.1 \times 10^{-3})^3$ ($= 3.88 \times 10^{-8} \text{ m}^3$)	C1
	$\rho = 4.8 \times 10^{-4} / 9.81 \times V$	C1
	$= 1300 \text{ kg m}^{-3}$	A1
1(c)(i)	W downwards, U upwards, F_v upwards	B1
1(c)(ii)	$F_v = 7.2 \times 10^{-4} - 4.8 \times 10^{-4}$ $= 2.4 \times 10^{-4} \text{ (N)}$	C1
	velocity $= 2.4 \times 10^{-4} / (17 \times 2.1 \times 10^{-3})$ $= 6.7 \times 10^{-3} \text{ m s}^{-1}$	A1

Question	Answer	Marks
2(a)	force (on droplet of water) in horizontal direction is zero.	B1
2(b)	(time taken \Rightarrow) $3.5 / 6.6 = 0.53 \text{ (s)}$	A1
2(c)	$s = ut + \frac{1}{2}at^2$ $s = \frac{1}{2} \times 9.81 \times 0.53^2$	C1
	$h = 1.4 \text{ m}$	A1

Question	Answer	Marks
2(d)	displacement is straight-line distance (from P to Q) so less (than distance along path) or displacement is the shortest distance (from P to Q).	B1
2(e)	$(\text{displacement})^2 = 3.5^2 + 1.4^2$	C1
	displacement = 3.8 m	A1

Question	Answer	Marks
3(a)	$(m =) \rho V$	C1
	$= 1.0 \times 10^3 \times 1.5 \times 10^{-4} \times 5.0 \times 1.6 = 1.2 \text{ (kg)}$	A1
3(b)(i)	$(\Delta)p = 1.2 \times 5.0$ $= 6.0 \text{ N s}$	A1
3(b)(ii)	$F = 6.0 / 1.6$ or $1.2 \times 5.0 / 1.6$ $= 3.8 \text{ N}$	A1
3(c)	Newton's third law applies (so) 3.8 N.	B1
3(d)	$p = F / A$ $= 3.8 / 1.5 \times 10^{-4}$	C1
	$= 2.5 \times 10^4 \text{ Pa}$	A1

Question	Answer	Marks
4(a)	ratio = $300 / 3200$ = 0.094	A1
4(b)	$E = \frac{1}{2}mv^2$ or $E \propto v^2$	C1
	ratio = $(0.094)^{0.5}$ = 0.31	A1
4(c)	work (done against frictional force) = $3200 - 300$ (=2900)	C1
	length = $2900 / 76$ = 38 m	A1
4(d)(i)	$E = \frac{1}{2}kx^2$ or $E = \frac{1}{2}Fx$ <u>and</u> $F = kx$	C1
	$140 = \frac{1}{2} \times 63 \times x^2$ or $140 = \frac{1}{2}Fx$ <u>and</u> $F = 63x$	
	$x = 2.1$ m	A1
4(d)(ii)	percentage efficiency = $(140 / 300) \times 100$ = 47%	A1
4(d)(iii)	curved line from the origin	M1
	gradient of line increases	A1

Question	Answer	Marks
5(a)(i)	(two) waves travelling (at same speed) in opposite directions overlap	B1
	waves (of the same type) have same frequency/wavelength	B1
5(a)(ii)	phase difference = 0	A1

PUBLISHED

Question	Answer	Marks
5(b)(i)	$f_0 = f_s v / (v - v_s)$ $543 = f \times 334 / (334 - 13)$	C1
	$f = 522 \text{ Hz}$	A1
5(b)(ii)	(the speed is) decreasing	B1
5(c)(i)	$I \propto A^2$	B1
	$I_T / I_0 = \cos^2 20^\circ$ or $A_T / A_0 = \cos 20^\circ$	C1
	ratio = 0.94	A1
5(c)(ii)	angle = 140°	A1

Question	Answer	Marks
6(a)(i)	$P = VI$	C1
	$I = 36 / 8.0$ $= 4.5 \text{ A}$	A1
6(a)(ii)	charge = 4.5×50 $= 225$	C1
	number = $225 / 1.6 \times 10^{-19}$ $= 1.4 \times 10^{21}$	A1
6(a)(iii)	$R = V^2 / P$ or $R = V / I$ or $R = P / I^2$ $= 8.0^2 / 36$ or $= 8.0 / 4.5$ or $= 36 / 4.5^2$	C1
	$= 1.8 \Omega$	A1

PUBLISHED

Question	Answer	Marks
6(b)	$R = \rho L / A$	C1
	$L = (1.8 \times 0.25 \times 10^{-6}) / 1.4 \times 10^{-6}$ = 0.32 m	A1
6(c)	(larger cross-sectional area, same length, same resistivity and so) less resistance	M1
	(same p.d. and more current so) more power (dissipated)	A1
6(d)	current (in wire) is the same	M1
	(same p.d. across wire so) power stays the same	A1

Question	Answer	Marks
7(a)(i)	(electron) neutrino	B1
7(a)(ii)	nucleon number = 22	A1
	proton number = 10	A1
7(a)(iii)	up up down changes to up down down or up changes to down	B1
7(b)(i)	charge = $-\frac{2}{3} e$	A1
7(b)(ii)	antiup / anticharm / antitop	B1